www.hep-druginteractions.org

Antiepileptic Treatment Selector

and use only. Not for distribution

r personal use only. Not for distrib

onai use only. Not jor alstri

Full information available at www.hep-druginteractions.org istribution. For personal use only. Not for distribution.

	DCV	ELB/GZR	G/P	LED/SOF	OBV/PTV/r	OBV/PTV/r +DSV	RDV	SOF	SOF/VEL	SOF/VEL/VOX
Carbamazepine	U a	Ų	↓ b, c	↓ь	∬ a	Ûа	₩	↓ b, e	↓ь	↓
Clobazam	\leftrightarrow	\leftrightarrow	\leftrightarrow	\leftrightarrow	↑f	↑f	\leftrightarrow	\leftrightarrow	\leftrightarrow	\leftrightarrow
Clonazepam	\leftrightarrow	\leftrightarrow	\leftrightarrow	\leftrightarrow	↑	1	\leftrightarrow	\leftrightarrow	\leftrightarrow	\leftrightarrow
Eslicarbazepine	∱a	₩	↓ь	\leftrightarrow	Ų	↓	\leftrightarrow	\leftrightarrow	ŲÞ	Ų
Diazepam	\leftrightarrow	\leftrightarrow	\leftrightarrow	\leftrightarrow	↑↓ ^h	↑↓ ^h	\leftrightarrow	\leftrightarrow	\leftrightarrow	\leftrightarrow
Ethosuximide	\leftrightarrow	\leftrightarrow	\leftrightarrow	\leftrightarrow	↑	↑	\leftrightarrow	\leftrightarrow	\leftrightarrow	\leftrightarrow
Gabapentin	\leftrightarrow									
Lacosamide	\leftrightarrow	\leftrightarrow	\leftrightarrow	\leftrightarrow	1	1	\leftrightarrow	\leftrightarrow	\leftrightarrow	\leftrightarrow
Lamotrigine	\leftrightarrow	\leftrightarrow	\leftrightarrow	\leftrightarrow	\downarrow	Ļ	\leftrightarrow	\leftrightarrow	\leftrightarrow	\leftrightarrow
Levetiracetam	\leftrightarrow									
Lorazepam	\leftrightarrow	\leftrightarrow	\leftrightarrow	\leftrightarrow	1	1	\leftrightarrow	\leftrightarrow	\leftrightarrow	\leftrightarrow
Midazolam (oral)	\leftrightarrow	Ť	\leftrightarrow	\leftrightarrow	1	↑ (\Rightarrow	\leftrightarrow	\leftrightarrow	\leftrightarrow
Midazolam (parental)	\leftrightarrow	Ť	\leftrightarrow	\leftrightarrow	↑ ⁱ	↑ ⁱ	\leftrightarrow	\leftrightarrow	\leftrightarrow	\leftrightarrow
Oxcarbazepine	∱a	⇒	↓ь	↓ь	₽	Ų	₽	₽₽	↓ь	₩
Perampanel	\leftrightarrow	\leftrightarrow	\leftrightarrow	\leftrightarrow	1	↑ (\Rightarrow	\leftrightarrow	\leftrightarrow	\leftrightarrow
Phenobarbital	Ų a	₽	↓ь	₽р	↓↓	↓↓	⇒	₽₽	∜ь	₩
Phenytoin	Ų a	Ĥ	₿₽	↓ь	↓↓	↓↓	₩	∯ь	Ų₽	ţ
Pregabalin	\leftrightarrow									
Primidone	Ų a	Ĥ	₽₽	₽	Ų↑	Ų↑	₩	₿₽	₿ъ	ţ
Retigabine	\leftrightarrow									
Rufinamide	↓	₽	₽	₩	Ų	Û	₽	₽	₩	Ų
Sultiame	\leftrightarrow	\leftrightarrow	\leftrightarrow	\leftrightarrow	∱ f	↑f	\leftrightarrow	\leftrightarrow	\leftrightarrow	\leftrightarrow
Tiagabine	\leftrightarrow	\leftrightarrow	\leftrightarrow	\leftrightarrow	↑f	↑f	\leftrightarrow	\leftrightarrow	\leftrightarrow	\leftrightarrow
Topiramate	\leftrightarrow									
Valproate semisodium (divalproex sodium)	\leftrightarrow	\leftrightarrow	\leftrightarrow	\leftrightarrow	ţ	↓i	\leftrightarrow	\leftrightarrow	\leftrightarrow	\leftrightarrow
Vigabatrin	\leftrightarrow									
Zonisamide	\leftrightarrow									

Colour Legend

No clinically significant interaction expected.

These drugs should not be coadministered.

Potential interaction which may require a dosage adjustment or close monitoring.

Potential interaction predicted to be of weak intensity.

Text Legend

- ↑ Potential increased exposure of the anticonvulsant
- Potential decreased exposure of the anticonvulsant
- Potential increased exposure of HCV DAA
 Potential decreased exposure of HCV DAA

↔ No significant effect

Numbers refer to increased or decreased AUC as observed in drug-drug interaction studies.

- a Coadministration is contraindicated. However, a published case series demonstrates that clinical cure (guided by TDM) may be achieved in patients where coadministration cannot be avoided.
- b Coadministration is not recommended. However, reports indicate that patients who remained on anti-epileptics during HCV DAA therapy achieved SVR.
- c Glecaprevir AUC decreased by 66%; pibrentasvir AUC decreased by 51%.
- d Coadministration with ombitasvir/paritaprevir/ritonavir + dasabuvir decreased the AUCs of ombitasvir, paritaprevir and dasabuvir by 31%, 70% and 70%, respectively.
- e Coadministration decreased sofosbuvir C_{max} and AUC by 48%. C_{max} of GS-331007 increased by 4%; AUC decreased by 1%.
- f Close monitoring is recommended for signs and symptoms of increased antiepileptic concentration.
- g If coadministration is necessary, the dose of daclatasvir should be increased to 90 mg once daily.
- h Coadministration with ombitasvir/paritaprevir/ritonavir + dasabuvir increased diazepam C_{max} by 18%, but decreased AUC by 22%; nordiazepam C_{max} increased by 10%, but AUC decreased by 40%. Monitor closely and adjust dose if indicated.
- i Coadministration should take place under close clinical monitoring with medical management in case of respiratory depression. Dose reduction should be considered.
- I The clinical significance of this is unclear. No a priori dose adjustment is required. Perform therapeutic drug monitoring and adjust dose if indicated.

Abbreviations:

DCV Daclatasvir ELB/GZR Elb RDV Ravidasvir SOF Sofosbu

ELB/GZR Elbasvir/Grazoprevir SOF Sofosbuvir G/P Glecaprevir/Pibrentasvir VEL Velpatasvir LED Ledipasvir VOX Voxilaprevir OBV/PTV/r +DSV Ombitasvir/Paritaprevir/Ritonavir +Dasabuvir

© Liverpool Drug Interactions Group, University of Liverpool, 3rd Floor William Henry Duncan Building, 6 West Derby Street, Liverpool, L7 8TX.

We aim to ensure that information is accurate and consistent with current knowledge and practice. However, the University of Liverpool and its servants or agents shall not be responsible or in any way liable for the continued currency of information in this publication whether arising from negligence or otherwise howsoever or for any consequences arising therefrom. The University of Liverpool expressly exclude liability for errors, omissions or inaccuracies to the fullest extent permitted by law.

